

# On an Alternative Cosmology

May 21, 1998

## Introduction

The Standard Cosmological Model faces numerous problems (see, for example, [1-3]). The present work is an attempt to formulate an alternative cosmological concept to resolve the problems on a basis of a set of physical hypotheses.

## 1 Hypotheses

The Grand Universe (GU) is an infinite multitude of Typical Universes (TUs). TUs have limited volumes and are chaotically and uniformly, on average, dispersed in a Grand Universe Space (GUS). Matter, in all forms, outside the TUs is a Grand Universe Background (GUB). Our Home Universe (HU) is an example of a TU consisting of stars, black holes (if any), and other matter called a Typical Universe Background (TUB). A Local Universe (LU) is the visible part of the HU. Observed Cosmic Background Radiation (CBR) is a typical component of TUB. A Local Grand Universe (LGU) is an imaginary part of the GU with a random sample of TUs.

### Basic hypotheses in the GU concept

1. The GU exists in the GUS, which is an infinite, flat 3-space. The GU is in a steady state. It is a uniform and isotropic, on average, universe. It has both baryon and electric symmetry. It has no absolute reference system (absolute "aether").
2. The intrinsic properties of GU matter are relativistic motion and self-recreation.
3. The GU exists in the form of matter and antimatter TUs, which evolve due to their interaction with the GUB.

The given hypotheses are in principle sufficient for a numerical simulation of GU matter behavior but seem to be redundant. In the author's opinion, a generalized matter transport theory must be developed for this purpose. Let us consider a statistical ensemble of GU matter under conditions formulated in hypothesis (1). Material objects are supposed to interact in direct collision and

by gravitational forces while in flight. Conventional physical processes must be taken into account. Then all GU properties stated in hypotheses (2) and (3) are expected to be deduced from the theory. In particular, a GU matter ensemble has to be characterized by a smooth, self-sustained mass distribution (equally for matter and antimatter) in a broad mass range, beginning with elementary free particles and ending with a TU itself. The distribution function must be one of the main goals of the theory. The problem is that any evolving TU, being a sub-ensemble, may appear to be more complex than the ensemble itself. Further, we take the above set of hypotheses for granted to conjecture qualitative physical consequences and to draw a cosmological picture in the framework of the GU concept, as we call it.

## 2 Relativistic Motion of GU Matter

The postulated absence of absolute aether in GUS means that a coordinate-momentum distribution of GU matter must be a Lorentz invariant. In other words, in any reference system it takes a form:

$$f(x_1, x_2, x_3; p_1, p_2, p_3) = \text{const} \quad (1)$$

This characterizes the chaotic motion of a relativistic photon or corpuscular gas [4]. In our case it applies to any material objects or gravitationally linked object systems as well. The integral over momentum distribution in (1) diverges because we deal with an ideal model of non-interacting gas. For interacting particles the distribution will have a smooth cut-off of upper energies in correspondence with limited energy density of GUB.

## 3 TU Origin and Evolution. The Expanding Universe.

Relativistic motion of GU matter is a cause of a steady state and cyclic GU recreation in the processes of annihilation and pair creation. One can imagine a probability chain of random object collisions resulting in a smooth mass distribution of separate bodies or gravitationally linked body systems. Small but certain chances are given to an initial matter (antimatter) coagulant to survive and grow to a "mature" TU age. We may call the relevant process a TU embryo evolution. As the TU evolves, it captures a large amount of cold matter from the GUB. For this reason a TU space must store thermal radiation emitted by the cold matter. Yet, cold matter provides for the origination of dust clouds and stars. Evolving stars converge some gravitational and nuclear energy into heat with a subsequent impact on TU properties. There is another reason for changing TU properties during its evolution. A TU of greater mass is able to capture bodies of higher speed. Eventually a TU becomes a relativistic system with growing instability. It cannot collapse but may easily decay at any stage

if its kinetic energy happens to exceed its potential. A rough criterion of decay may be written as:

$$\frac{GM}{Rc^2(\gamma - 1)} < 1 \quad (2)$$

where:  $M$ -mass,  $R$ -mean radius,  $\gamma$ -effective Lorentz factor,  $G$ -gravitational constant,  $c$ -speed of light.

Thus, a TU is evolving due to its active interaction with GUB but inherent factors are essential. An energy distribution of TU matter is different from that of GUB. First, the mean energy is smaller. Second, there must be a space-energy correlation: high speed objects move with higher radii.

Keeping in mind this picture of TU evolution, let us consider our HU. Tracing back to red shifts,  $z \sim 10$ , we can imagine the HU being about three orders more dense than at present, having a mass of roughly  $M \sim 10^{52} \text{kg}$  and a radius  $R \sim 4 * 10^{25} \text{m}$ , which is apparently close to criterion (2). The most effective mechanism triggering the HU to decay might be an abrupt mass drop due to annihilation in a process of accidental "soft collision" with a smaller antimatter TU. As a result, we observe the present Expanding Universe with the Hubble's expansion law. Some features of the Expanding Universe are discussed below.

## 4 Quasars

We have to explain several quasar features. An effective volume of a quasar "engine" is about  $10^{37} \text{m}^3$  with a power density of about  $10^3 \text{Wm}^{-3}$ . Quasars came into existence in the last epoch of maximal  $z$  during what is thought to be the start of expansion. Before that, a long period of HU pre-expansion evolution is supposed to have taken place. Quasars may vary their power by about 50% in a short period of time. Finally, they are apparently located in the center of galaxies. One can start by recalling that annihilation is a process of potentially the highest power density possible in conventional physics. We can assess  $3 * 10^9$  nucleon pair annihilation in a unit volume per second for a power rate of  $1 \text{Wm}^{-3}$ . Suppose an "engine" volume is filled in with rock-like particles of a size  $\alpha$  and density  $\rho$ . Let particle clouds be mixed with corpuscular antimatter radiation with a flux  $\varphi$ . Then an annihilation rate,  $A$ , is:

$$A = \rho \alpha^2 \varphi \quad (3)$$

Assuming  $\rho = 10^{-6} \text{m}^{-3}$ ,  $\alpha = 10^{-1} \text{m}$ ,  $\varphi = 3 * 10^{20} \text{m}^{-2} \text{s}^{-1}$  we have the required quasar power. This rough estimate shows a physical feasibility of the annihilation nature of quasars. Presumably they are active gravitating centers where exploding processes cause varying gas pressure and subsequently varying power.

Gamma rays bursts having cosmological origin have been discovered some years ago. We think the phenomenon has the same physical nature as quasars. Antimatter objects constantly penetrating from the GUB cause random annihilation pulses when colliding with HU stars. If so, a statistical analysis of gamma rays bursts should reveal a HU boundary.

## 5 Redshift Distribution

We assume a physical space to be Minkovski's space. Its space-time scales are proportional to those of a conformally flat space taken in the original Freedman's solution for an expanding universe. For simplicity, we neglect a peculiar velocity field and take a model of a uniformly expanding coherent gas of particles. A gas system in a form of an expanding sphere has an initial effective radius  $R(0)$ . In this case the special relativity theory gives us a velocity distribution of moving particles [5]:

$$f(\beta)d\beta = Const * \gamma^4 \beta^2 d\beta \quad (4)$$

for:  $0 \leq \beta \leq \beta_{max}$  where:  $\beta_{max}$  is determined by  $R(0)$ .

As usual,  $\gamma = (1 - \beta^2)^{-1/2}$ .

A velocity distribution (4) corresponds to a momentum distribution (1) if one takes into account that relativistic mass depends on velocity. Hence, an initial chaotic distribution is consistent with a uniform HU expansion. Neglecting a peculiar velocity field means that initial velocities are supposed to have radial alignment.

If we take a velocity distribution (4) as an approximate picture seen by a imaginary observer at the HU center and at the initial moment,  $t = 0$ , we can easily find a corresponding redshift distribution  $\varphi_i(z)$ :

$$\varphi_i(z) = f(\beta(z)) \frac{d\beta}{dz} \quad (5)$$

with a special relativity relation:

$$\beta(z) = \frac{(1+z)^2 - 1}{(1+z)^2 + 1} \quad (6)$$

Then:

$$\varphi_i(z) = const * \frac{z^2(2+z)^2}{4(1+z)^3} \quad (7)$$

In a process of expansion a deceleration influences velocity and corresponding redshift distributions making them dependent on time. A radial coordinate of some luminous, moving object is:

$$r(t) = c \int_0^t \beta(t) dt \quad (8)$$

Suppose a light packet was emitted at emission moment  $t_e$  and registered at the moment of observation  $t_0$ . Then an object velocity at  $t = t_e$  is found from the equation:

$$\int_0^{t_e} \beta(t) dt = (t_0 - t_e) \quad (9)$$

Therefore, a set of  $t_e$  values and corresponding  $\beta_e$  values is available for finding an emission velocity distribution  $\varphi_0(\beta_e)$  and a corresponding redshift distribution  $\varphi_0(z)$  observed at  $t = t_0$ .

$$\varphi_0(\beta_e) = \varphi_i(\beta_i(\beta_e)) \frac{d\beta_i}{d\beta_e} \quad (10)$$

and

$$\varphi_0(z) = \varphi_0(\beta_e(z)) \frac{d\beta}{dz} \quad (11)$$

for:  $0 \leq z \leq z_{max}(\beta_{e_{max}})$

The last expression concludes the solution of the redshift problem. At small  $z$ :

$$\varphi_0(z) = z^2 \quad (12)$$

and, the smaller the  $\beta_{e_{max}}$ , the broader the range. During the present epoch redshift counts show the dependence (12) at least up to  $z \approx 0.25$  [1]. This is evidence that appreciable time passed since the HU started expanding. Detailed numerical analysis in the GU concept framework must reveal a good deal of knowledge about our HU.

The Standard Model failed to predict redshift counts. The reason for this failure, as well as criticism of the Standard Model, is not a subject of this work.

## 6 “Black Matter” and CBR

As discussed in previous sections, a TU captures cold matter from the GUB or creates its own in a form of star remnants during its evolution. This is baryon cold matter, referred to in the Standard Model as ”Black Matter”. It is uniformly distributed, on average, in TU space. Cosmic background radiation (CBR) is a thermal radiation partially in relaxation with cold matter. The surface temperature of cold matter in the HU is 2.7 K at present. In general this temperature is determined by the local energy balance. Apparently, a balance of absorbed and emitted energy is negative, and cold matter is cooling in a process of expansion. Its temperature seems to be inversely proportional to an expansion factor. A detailed study of the relationship between CBR and cold matter in the GU concept framework must clarify the situation.

## 7 Cosmic Rays

The GU concept gives a natural explanation of high energy cosmic rays observed from Earth: an ultrarelativistic GUB gas must be their source. Therefore, cosmic rays are a result of GUB gas transformation and moderation in a process of transport through an HU medium. Qualitatively we may assume a logarithmic energy loss being proportional to a time of GUB particle travel in an HU medium:

$$\frac{d\varepsilon}{\varepsilon} = -\alpha dt \quad (13)$$

and a Poisson probability to reach an observer:

$$w(t)dt = \frac{dt}{T} \exp(-\frac{t}{T}) \quad (14)$$

where  $\alpha$ ,  $T$  are transport model parameters.

Then an observed spectrum is:

$$F_0(\varepsilon) = Const * \varepsilon^{-(1+1/\alpha T)} \quad (15)$$

which is formed in a broad energy range  $\varepsilon < \varepsilon_i$  ( $\varepsilon_i$  is the initial energy of relativistic GUB particles). For  $\alpha T = 2/3$  (which is a logarithmic energy loss per effective particle life time) we have a spectrum close to that observed.

A GUB gas is baryon symmetric but most primary antiparticles must have been annihilated in the transport process, excepting those in the upper energy range. Unfortunately, measurement and identification of cosmic rays at super-high energies is an extremely difficult task.

Both CBR and cosmic rays are part of a TUB, therefore they play a role of absolute aether participating in an expansion process. However, boundary effects must cause some departures from Hubble's law. This means that an observer who is not in the center of the HU must see an asymmetry in angular distribution. Some anisotropic effects are really found, but higher measurement precision is needed for their interpretation.

## 8 Some Other Cosmological Issues

There are Standard Model problems of singularity, flatness, horizon, and "universal

aether" which have a quite general and, to some extent, philosophical character. Fortunately, these do not arise in the alternative cosmology concept. Other specific physical problems are related to the common question of whether Hubble's time is long enough comparing with the time required for star evolution and recycling, galaxy formation and clustering, large-scale structuring, nucleosynthesis, and so on. The alternative concept does not impose time constraints on this sort of problems because all processes are considered in a time scale that includes the GU recreation cycle and the pre-expansion stage of the HU.

The alternative concept predicts physical boundary of any TU, the HU in particular. Therefore, the HU must have a rotation motion in a GUS, and there must be some evidence of it. Subsequent anisotropic effects in the HU picture will depend on our location.

## Conclusion

The concept qualitatively explains available observational data for our Home Universe but predictions concerning the whole Universe seem to have the same level of likelihood. A general transport theory is needed to investigate in detail the suggested cosmological framework.

## Acknowledgements

I would like to thank Prof. N. Rabotnov for fruitful discussion of the problem. I also wish to acknowledge Prof. J. Peebles for encouraging contacts.

## References

1. P.J.E. Peebles. *Principles of Physical Cosmology*. Princeton University Press. 1993.
2. *Critical Dialogues in Cosmology*. In celebration of the 250th anniversary of Princeton University. 24-27 June, (1996). World Scientific Press.
3. M. Rees. *Before the Beginning*. Helix Books, Addison-Wesley Press, 1997.
4. J.L. Synge. *The Relativistic Gas*. North-Holland Publishing Company, Amsterdam. 1997.
5. V.A. Fok. *Theory of Space, Time, and Gravity*. (in Russian) Published in Moscow. Teoriya prostranstva, vremeni i tyagoteniya. Gosizdatelstvo tekhniko-teoreticheskoi literaturi. Moskva, 1955.